



Claims

1. A process for producing an optically active 3-(4-hydroxyphenyl)propionic acid of the formula (6):



wherein R^2 is an alkyl group, R^5 to R^8 are each independently a hydrogen atom or a substituent; and the symbol * is a chiral carbon atom,

or a salt thereof, which comprises reacting a benzaldehyde of
10 the formula (1):




wherein R^1 is a protective group; and R^5 to R^8 are each the same as defined above,

with a glycolic acid derivative of the formula (2):




wherein R^3 is a hydrocarbon group, and R^2 is the same as defined above,

hydrolyzing the resulting product to give a cinnamic acid of the formula (4):




wherein R^1 , R^2 , and R^5 to R^8 are each the same as defined above,
 or a salt thereof, and subjecting the cinnamic acid (4) or a
 salt thereof to asymmetric hydrogenation to give an optically
 5 active phenylpropionic acid of the formula (5):




wherein all the symbols are each the same as defined above,
 or a salt thereof, followed by deprotection.

10 2. A process for producing an optically active 3-(4-
 hydroxyphenyl)propionic acid of the formula (6):



15 wherein R^2 is an alkyl group; R^5 to R^8 are each independently
 a hydrogen atom or a substituent; and the symbol * is a chiral
 carbon atom,
 or a salt thereof, which comprises reacting a benzaldehyde of
 the formula (1):




wherein R^1 is a protective group; and R^5 to R^8 are each the same as defined above,
with a glycolic acid derivative of the formula (2):




5

wherein R^3 is a hydrocarbon group, and R^2 is the same as defined above, followed by hydrolysis to give a cinnamic acid of the formula (4):




10 wherein R^1 , R^2 , and R^5 to R^8 are each the same as defined above,
or a salt thereof, and subjecting the cinnamic acid (4) or a
salt thereof to asymmetric hydrogenation.

3. A process for producing an optically active 3-(4-
15 hydroxyphenyl)propionic acid of the formula (6):



wherein R^2 is an alkyl group; R^5 to R^8 are each independently a hydrogen atom or a substituent; and the symbol * is a chiral carbon atom,


or a salt thereof, which comprises reacting a
5 4-hydroxybenzaldehyde of the formula (7):



wherein R^5 to R^8 are each the same as defined above, with a glycolic acid derivative of the formula (2):



10 wherein R^3 is a hydrocarbon group; and R^2 is the same as defined above, followed by hydrolysis to give a 4-hydroxycinnamic acid of the formula (9):




wherein R^2 , and R^5 to R^8 are each the same as defined above,
15 or a salt thereof, and subjecting the 4-hydroxycinnamic acid (9) or a salt thereof to asymmetric hydrogenation.

4. The process according to any one of claims 1 to 3,
wherein the asymmetric hydrogenation is carried out in the
20 presence of a chiral catalyst.


5. The process according to any one of claims 1 to 4, wherein the chiral catalyst is a transition metal complex.

5 6. The process according to claim 5, wherein the transition metal complex is a complex of the metal of Groups 8 to 10 in the periodic table.

7. A process for producing an optically active
10 carboxylic acid of the formula (12):



wherein R^{11} and R^{12} are each independently a hydrogen atom or a substituent; R^{13} is a hydrogen atom, an optionally substituted hydrocarbon group or a metal atom; R^{14} is a hydrogen atom or
15 a protective group; and the symbol * is an chiral carbon atom, or a salt thereof, which comprises subjecting an α,β -unsaturated carboxylic acid of the formula (11):



wherein R^{11} to R^{14} are each the same as defined above,
20 or a salt thereof, to asymmetric hydrogenation in the presence of a transition metal complex, provided that when the transition metal complex is rhodium, the protective group represented by R^{14} in the above formula (11) is a group other than acyl.

8. The process according to claim 7, wherein the transition metal complex is a complex of the metal of Groups 8 to 10 in the periodic table.

5

9. The process according to claim 1 or 3, wherein the chiral catalyst is a mixture of a chiral ligand and a transition metal compound.

10

10. The process according to any one of claims 1 to 3, wherein the optically active phenylpropionic acid of the formula (5) or a salt thereof obtained by the method according to any one of claims 1 to 3 is crystallized from a solvent.

15

11. The process according to claim 10, wherein the solvent used for the crystallization is a member selected from the group consisting of hydrocarbons, alcohols, ketones and water, and a mixture thereof.


20

12. The process according to any one of claims 1 to 3, wherein the optically active 3-(4-hydroxyphenyl)propionic acid of the formula (6) or a salt thereof obtained by the method according to any one of claims 1 to 3 is crystallized from a solvent.

25


13. The process according to claim 12, wherein the solvent used for the crystallization is a member selected from the group consisting of aromatic hydrocarbons, aliphatic hydrocarbons, alcohols and water, and a mixture thereof.

14. A process for producing an optically active phenylpropionic acid of the formula (5):



5 wherein R¹ is a protective group; R² is an alkyl group; R⁵ to R⁸ are each independently a hydrogen atom or a substituent; and the symbol * is an chiral carbon atom,
or a salt thereof


which comprises subjecting a cinnamic acid of the formula (4):



10


wherein R¹, R², and R⁵ to R⁸ are each the same as defined above,
or a salt thereof,
to asymmetric hydrogenation.

15 15. A process for producing an optically active 3-(4-hydroxyphenyl)propionic acid of the formula (6):




wherein R^2 is an alkyl group; R^5 to R^8 are each independently a hydrogen atom or a substituent; and the symbol * is a chiral carbon atom,

or a salt thereof, which comprises subjecting a cinnamic acid
5 of the formula (4):



wherein R^1 , R^2 , and R^5 to R^8 are each the same as defined above, or a salt thereof, to asymmetric hydrogenation.


10 16. A process for producing an optically active 3-(4-hydroxyphenyl)propionic acid of the formula (6):



15 wherein R^2 is an alkyl group; R^5 to R^8 are each independently a hydrogen atom or a substituent; and the symbol * is a chiral carbon atom,


or a salt thereof,

which comprises subjecting a 4-hydroxycinnamic acid of the formula (9):




wherein R^2 , and R^5 to R^8 are each the same as defined above, or a salt thereof to asymmetric hydrogenation.

- 5 17. A process for producing an optically active 3-(4-hydroxyphenyl)propionic acid of the formula (6):




wherein R^2 is an alkyl group; R^5 to R^8 are each independently a hydrogen atom or a substituent; and the symbol * is a chiral carbon atom,

10 or a salt thereof, and an optically active phenylpropionic acid of the formula (5):




wherein R^1 is a protective group; and R^2 , R^5 to R^8 and the symbol *

15 * are each the same as defined above, or a salt thereof, which comprises subjecting a cinnamic acid of the formula (4):




wherein R^1 , R^2 , and R^5 to R^8 are each the same as defined above, or a salt thereof, to asymmetric hydrogenation.

- 5 18. A process for producing an optically active 3-(4-hydroxyphenyl)propionic acid of the formula (6):



wherein R^2 is an alkyl group, R^5 to R^8 are each independently a hydrogen atom or a substituent; and the symbol * is a chiral carbon atom,

10 or a salt thereof, which comprises reacting a benzaldehyde of the formula (1):



wherein R^1 is a protective group; and R^5 to R^8 are each the same


15 as defined above,

with a glycolic acid derivative of the formula (2):




wherein R^3 is a hydrocarbon group, and R^2 is the same as defined above,


hydrolyzing the resulting product to give a cinnamic acid of
5 the formula (4):



wherein R^1 , R^2 , and R^5 to R^8 are each the same as defined above,
or a salt thereof, and subjecting the cinnamic acid (4) or a
salt thereof to asymmetric hydrogenation to give an optically
10 active phenylpropionic acid of the formula (5):



wherein all the symbols are each the same as defined above,
or a salt thereof, and an optically active
3-(4-hydroxyphenyl)propionic acid of the formula (6):




15

wherein all the symbols are each the same as defined above,
or a salt thereof, followed by deprotection.

Abstract

The present invention relates to a process for producing an optically active 3-(4-hydroxyphenyl)propionic acid useful
 5 as intermediates for medicines, through short steps in good yield and with high optical purity. More specifically, the present invention relates to a process for producing an optically active 3-(4-hydroxyphenyl)propionic acid of the formula (6):




10

wherein R^2 is an alkyl group; R^5 to R^8 are each independently a hydrogen atom or a substituent; and the symbol * is an chiral carbon atom,

15

or a salt thereof, which comprises reacting a benzaldehyde of the formula (1):



wherein R^1 is a protective group; and R^5 to R^8 are each the same as defined above,

with a glycolic acid derivative of the formula (2):




20


wherein R^3 is a hydrocarbon group; and R^2 is the same as defined

above,

hydrolyzing the resulting product to give a cinnamic acid of the formula (4):



- 5 wherein R¹, R² and R⁵ to R⁸ are each the same as defined above, or a salt thereof, and subjecting the resulting cinnamic acid (4) or a salt thereof to asymmetric hydrogenation to give an optically active phenylpropionic acid of the formula (5):



- 10 wherein all the symbols are each the same as defined above, or a salt thereof, followed by deprotection.